

Defusing COVID-19: Lessons Learned from a Century of Pandemics

Graciela Mujica ¹, Zane Sternberg ², Jamie Solis ¹, Taylor Wand ¹, Peter Carrasco ^{3,4}, Andrés F. Henao-Martínez ⁵ and Carlos Franco-Paredes ^{5,6,*}

¹ [University of Colorado School of Medicine](#), 13001 E 17th Pl, Aurora, CO 80045, USA; graciela.mujica@cuanschutz.edu (G.M.); jamie.solis@cuanschutz.edu (J.S.); taylor.wand@cuanschutz.edu (T.W.), ² [University of Colorado Denver](#), 13001 E 17th Pl, Aurora, CO 80045 USA; zane.sternberg@cuanschutz.edu ³ [International Association for Immunization Managers](#), Washington, DC, 20037 USA; rcphs@aol.com ⁴ Department of Immunization and Vaccines, World Health Organization, [Geneva](#), Switzerland ⁵ Division of Infectious Diseases, Department of Medicine, Anschutz Medical Center, University of Colorado, Aurora, CO 80220, USA; andres.henaomartinez@cuanschutz.edu ⁶ Hospital Infantil de México, Federico Gomez, [México City](#), 06720 México * Correspondence: Carlos.franco-paredes@cuanschutz.edu

BACKGROUND

- There are common patterns to the stories told as pandemics unfold.
- Amidst COVID-19, we found it crucial to identify and apply lessons from the behavior and spread of previous pandemics to gain insight into their interruption.
- We conducted a historical review of pandemics in the 20th century and compared their epidemiological aspects, as well as their social, environmental, and public health impacts.
- Our goal was to understand the biosocial influences on the rise and spread of pandemics to identify the most effective public health responses.


Pandemic	Total Deaths (Globally)	Case Fatality Rate (%)	Basic Reproductive Number (R ₀)
Spanish Flu (1918–1919)	20–50 mill	CMR ~2.5% *	2.11–2.5
Asian Flu (1957)	700,000–1.5 mil	0.02–0.05%	1.8
Hong Kong Flu (1968)	500,000–2.0 mil	0.67%	1.28–1.58
Swine Flu (2009)	2185–284,000	1.09%	1.33–1.38
SARS (2003)	774	9.6%	2.4
MERS (2012)	858	34.4%	1.1–1.2
COVID-19 (2020)	1,385,000	2–3%	1.5–3.5

Table 1. * Crude mortality rate (CMR) of ~2.5% calculated based on excess mortality observations

METHODS

- Obtained epidemiological data on coronaviruses and influenza viruses from the U.S. Public Health Service, World Health Organization (WHO), and peer-reviewed publications.
- Analyzed the relationship between peak pandemic death rates, time of the year, and population characteristics (“biosocial” drivers of pandemic spread).
- Assessed various non-pharmacologic interventions (NPIs), the timing of their implementation, and the regional impact these had on subsequent pandemic waves.

RESULTS



Virus Pandemic Years	Waves (Duration)	Timing
H3N8 - Russian Flu 1889–1890	3 (3 years)	W1: 1889–1890 / W2: 1890–1891 / W3: 1892–1893
H1N1 - Spanish Flu 1918–1919	3 (9 months)	W1: Jun–Jul 1918 / W2: Oct–Nov 1918 / W3: Feb–Mar 1919
H2N2 - Asian Flu 1957–1959	3 (15–18 months)	W1: Sep–Dec 1957 / W2: Dec–Mar 1958 / W3: Dec 58–Apr 59
H3N2 - Hong Kong 1968–1970	2 (12–18 months)	Based on global circ. / W1: Jul 68 to Aug 69 / W2: Sept 69–Sept 70
H1N1 - Swine Flu 2009–2010	1 Epidemic, 2 Pandemics (12 months)	E1: Jan–Mar 2009 / W1: Mar–Aug 2009 / W2: Aug 09–Jan 10

Table 2. Comparison of duration and wave seasonality of influenza pandemics from late 19th to 21st century

REFERENCES

- McNeill, W.H.; McNeill, W. *Plagues and Peoples*; Anchor: New York, NY, USA, 1998.
- Morens, D.M.; Daszak, P.; Markel, H.; Taubenberger, J.K. Pandemic COVID-19 Joins History’s Pandemic Legion. *mBio* **2020**, *11*, doi:10.1128/mbio.00812-20.
- World Health Organization. Report of the WHO-China Joint Mission on Coronavirus Disease 2019 (COVID-19). 2020.
- Johns Hopkins Coronavirus Resource Center. Available online: <https://coronavirus.jhu.edu/map.html> (accessed on 30 October 2020).
- Mummert, A.; Weiss, H.; Long, L.-P.; Amigó, J.M.; Wan, X.-F. A Perspective on Multiple Waves of Influenza Pandemics. *PLoS ONE* **2013**, *8*, e60343, doi:10.1371/journal.pone.0060343.
- Wang, L.-F.; Shi, Z.; Zhang, S.; Field, H.; Daszak, P.; Eaton, B.T. Review of Bats and SARS. *Emerg. Infect. Dis.* **2006**, *12*, 1834–1840, doi:10.3201/eid1212.060401.
- Kilbourne, E.D. Influenza Pandemics of the 20th Century. *Emerg. Infect. Dis.* **2006**, *12*, 9–14, doi:10.3201/eid1201.051254.
- Viboud, C.; Grais, R.F.; Lafont, A.B.; Miller, M.A.; Simonsen, L. Multinational Influenza Seasonal Mortality Study Group: Multinational impact of the 1968 Hong Kong influenza pandemic: Evidence for a smoldering pandemic. *Int. J. Infect. Dis.* **2005**, *192*, 233–248.

CONCLUSIONS

- In the past 100 years, the majority of pandemics have been caused by respiratory viruses that emerged during the winter in the Northern Hemisphere, during the second half of the typical flu season.
- All pandemics have originated from a non-human source. Most have exhibited the ability to cross from humans back into certain animal populations, thus serving as a site for additional antigenic shifts or genetic recombination events. Given that SARS-CoV-2 is capable of infecting animal hosts, similar events could result in viral mutations that impact herd immunity.
- The first 2–3 waves of a pandemic are most likely to occur within 12–15 months of the virus originating. That being said, the speed at which pandemics spread has increased significantly given the rise in population density, globalization, and ease of travel. Pandemics are becoming more common and more difficult to control.
- School-age children have been major drivers of household transmission and pandemic spread. Early institution of mitigation interventions such as school closures, interruption of mass gathering, travel restrictions, and community quarantine is critical to achieving at least a 50% reduction in transmission, which correlates with a basic reproductive number lower or equal to one ($R_0 \leq 1$). These have historically had the largest, most cost-effective impacts on incidence and mortality.
- Pandemics disproportionately impact socially disadvantaged populations. They do not produce inequalities but reveal issues we have failed to address as mandates of public health.

ACKNOWLEDGEMENTS

Authors’ Contributions: G.M., Z.S., C.F.-P., J.S. drafted the initial manuscript and reviewed the literature. Edited and approved the last version. A.F.H.-M., P.C., T.W. reviewed the literature, edited, and approved the last version. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Conflicts of Interest: The authors declare no conflict of interest.