

Return to Duty as a Functional Outcome Measure Following Orthopaedic Surgery: A Preliminary Investigation

F. James Powlan – University of Colorado School of Medicine

Introduction

In the military, return to duty (RTD) status has commonly been used as a functional outcome measure following orthopaedic surgery. This is at times regarded similarly to “return to sport (RTS),” or as an indicator of return to full function. The purpose of this study is to determine the efficacy of RTD status as a standalone surrogate for RTS and assess its overall usefulness as a functional outcome measure. The purpose of this study was to answer the following questions:

- (1) For military patients, is return to duty status an effective surrogate for return to sport following orthopaedic surgery?
- (2) Is return to duty status an indicator of returning to full function after orthopaedic surgery?

Methods

Pre and post-operative self-reported RTD status, profile status, RTS status, deployment status, MOS (military occupation specialty) changes, and MEB (medical evaluation board) status, were retrospectively reviewed for all active duty soldiers who underwent orthopaedic surgery at our institution from February 2017 to October 2018. Minimum follow up time for inclusion was 1 year.

Acknowledgements

B. Holt Zalneraitis MD, Nicholas Drayer MD, Matthew Nowak DO, Kyle Ardavanis MD, Brendan Masini MD, Daniel Kang MD

Results

Pre-Operative Profile Status

Status	%	n (140)
Full active duty with no restrictions	62.8	88
Profile for same joint/body part	23.6	33
Profile for different joint/body part	13.9	19

One-year Follow up Profile Status

Status	%	n (121)*
Full active duty with no restrictions	67.8	82
Profile for same joint/body part	24.8	30
Profile for different joint/body part	7.43	9
*Patients with pre-operative profile for different joint excluded		

One-year Follow up Modifiers

Modifiers	%	n (82)*
Sport-related questions		
Lower sport activity level	70.7	58
Limited in sport activity	12.2	10
Lower work level	29.3	24
Military specific modifiers		
Non-deployable	34.1	28
MEB	19.5	16
MOS change	3.66	3
*Patients reporting full active duty with no restrictions at one-year follow up (MEB = Medical Evaluation Board, MOS = Military Occupational Specialty)		

Literature Cited

Antosh JJ, Cameron KL, Marsh NA, Posner MA, DeBerardino TM, Svoboda SJ, Owens BD. Likelihood of Return to Duty Is Low After Meniscal Allograft Transplantation in an Active-duty Military Population. Clin Orthop Relat Res. 2020;478:722-730.
Antosh JJ, Patzkowski JC, Racusin AW, Aden JK, Waterman SM. Return to Military Duty After Anterior Cruciate Ligament Reconstruction. Mil Med. 2018;183:e83-e89.
Arroyo W, Misenheimer J, Cotter EJ, Wang KC, Fleida K, Pallas MP, Waterman BR. Effect of Anterior Anchor on Clinical Outcomes of Type II SLAP Repairs in an Active Population. Orthopedics. 2019;42:e32-e38.
Barrow AE, Sheean AJ, Burns TC. Return to duty following combat-related multi-ligamentous knee injury. Injury. 2017;48:861-865.
Burgmeier RJ, Hsu WK. Spine surgery in athletes with low back pain: considerations for management and treatment. Asian J Sports Med. 2014;5:e24284.
Cassidy JT, Hurley EJ, Moore D, Pauzenberger L, Mullett H. The majority of patients return to athletic activity following biceps tenodesis. Knee Surg Sports Traumatol Arthrosc. 2020.
Copp T. 126,000 service members in crosshairs for separation as DoD's 'deploy or get out' policy takes effect. Military Times.
Han PE, Zhang ZL, Chen CL, Han YC, Wei XC, Li PC. Comparison of primary arthrodesis versus open reduction with internal fixation for Lisfranc injuries: Systematic review and meta-analysis. J Postgrad Med. 2019;65:93-100.

Conclusions

RTD status is commonly reported in military orthopaedics as a way of describing post-operative functional outcome.

58.6% of patients reported full RTD with no restrictions/profile at final follow up. However, the majority of these patients reported lower level of sport activity.

Numerous patients also reported not returning to the same work level, being non-deployable, or undergoing an active MEB.

Therefore, our preliminary investigation suggests self-reported full RTD may not be an appropriate indicator of return to full function, nor an adequate surrogate for RTS.

It should be noted that our results are self reported as well as a lack of specific outcome scores for each surgery. Going forward using outcome scores based on surgery could give a more clear picture of RTD as a functional outcome measure.

Further Information

No financial disclosures nor conflicts of interest
Abstract presented at SOMOS 2020
Transcript submitted to *Clinical Orthopaedics and Related Research* January 2021
Email is franklin.powlan@cuanschutz.edu for further questions